



# STANDARDIZED ONE PAGE PHARMACY PRIOR AUTHORIZATION FORM

Mississippi Division of Medicaid, Pharmacy Prior Authorization Unit, 550 High St., Suite 1000, Jackson, MS 39201

Magnolia Health/Envolv Pharmacy Solutions  
Fax to: 1-877-386-4695 Ph: 1-866-399-0928  
<https://www.magnoliahealthplan.com/providers/pharmacy.html>

UnitedHealthcare/OptumRx  
Fax to: 1-866-940-7328 Ph: 1-800-310-6826  
<http://www.uhcommunityplan.com/health-professionals/ms/pharmacy-program.html>

Molina Healthcare/CVS Caremark  
Fax to: 1-844-312-6371 Ph: 1-844-826-4335  
<http://www.molinahealthcare.com/providers/ms/medicaid/pages/home.aspx>

Medicaid Fee for Service/Change Healthcare  
Fax to: 1-877-537-0720 Ph: 1-877-537-0722  
<https://medicaid.ms.gov/providers/pharmacy/pharmacy-prior-authorization/>

| BENEFICIARY INFORMATION   |  |
|---|--|
| Beneficiary ID: _____ - _____ - _____   | DOB: ____/____/_____   |
| Beneficiary Full Name: _____  |  |
| PRESCRIBER INFORMATION  |  |
| Prescriber's NPI: _____   |  |
| Prescriber's Full Name: _____   | Phone: _____   |
| Prescriber's Address: _____   | FAX: _____   |
| PHARMACY INFORMATION  |  |
| Pharmacy NPI: _____   |  |
| Pharmacy Name: _____  |  |
| Pharmacy Phone: _____   | Pharmacy FAX: _____  |
| CLINICAL INFORMATION  |  |
| Requested PA Start Date: _____ Requested PA End Date: _____   |  |
| Drug/Product Requested: _____ Strength: _____ Quantity: _____   |  |
| Days Supply: _____ RX Refills: _____ Diagnosis or ICD-10 Code(s): _____   |  |
| <input type="checkbox"/> Hospital Discharge   | <input type="checkbox"/> Additional Medical Justification Attached |
| Medications received through coupons and/or samples are not acceptable as justification.  |  |
| <b>PLEASE COMPLETE AND FAX DRUG SPECIFIC CRITERIA/ADDITIONAL DOCUMENTATION FORM FOUND BELOW</b>   |  |
| <i>Prescribing provider's signature (signature and date stamps, or the signature of anyone other than the provider, are not acceptable)</i> |  |
| I certify that all information provided is accurate and appropriately documented in the patient's medical chart.                            |  |
| Signature required: _____   | Date: _____  |
| Printed name of prescribing provider: _____   |  |

## FAX THIS PAGE

# CRITERIA/ADDITIONAL DOCUMENTATION

## HEPATITIS C

**FAX THIS PAGE**



### BENEFICIARY INFORMATION

Beneficiary ID: \_\_\_\_\_ - \_\_\_\_\_ - \_\_\_\_\_ DOB: \_\_\_\_/\_\_\_\_/\_\_\_\_

Beneficiary Full Name: \_\_\_\_\_

### Hepatitis C Therapy PA Request

**Diagnosis / Treatment Status (check all that apply) \*See Hepatitis-C PA description sheet for approval criteria and intolerance definitions.**

- Prescriber is, or has consulted with a gastroenterologist, hepatologist, ID specialist or other hepatic specialist. Requires consult within the past year with documentation of recommended regimen.
- Active HCV infection verified by viral load within the last year: HCV RNA: \_\_\_\_\_ million IU/mL Date: \_\_\_\_\_
- Genotype verified by lab:  1a  1b  2  3  4  5  6

HIV status:  positive  negative (required)

- Patient has not taken amiodarone within 535 days (required if regimen includes Harvoni or Sovaldi)

RBV-Ineligible reason: \_\_\_\_\_

Hepatic fibrosis stage \_\_\_\_\_

Last stage evaluation date: \_\_\_\_\_

Method of cirrhosis/fibrosis stage: \_\_\_\_\_

- Decompensated cirrhosis
- Compensated cirrhosis Child-Pugh Score and Date: \_\_\_\_\_
- Post-liver transplant Date: \_\_\_\_\_
- Hepatocellular carcinoma and awaiting a liver transplant: Transplant date: \_\_\_\_\_
  - Not yet scheduled
- Dialysis \_\_Yes/\_\_\_No
- CrCl \_\_\_\_ mL/min Lab Date w/n last year: \_\_\_\_\_
- Screened for HEP-B and HIV prior to HEP-C treatment start
  - Date of last test: Hep B: \_\_\_\_/\_\_\_\_/\_\_\_\_ HIV: \_\_\_\_/\_\_\_\_/\_\_\_\_
    - Timing of the screening for Hep-B/HIV should be based on patient specific risk factors but lab result date must be provided and if > 1 year ago, it should be documented in the record as to why repeat testing is not clinically warranted. **If the patient has had ongoing risk factors of any type, consider retesting in the month prior to HCV therapy. If positive, treatment must be considered per AASLD/IDSA and current NIH HIV guidelines.**
    - Repeat screening should be patient specific.

Patient is:  Treatment naïve  Relapser

If Relapser, then prior HCV Treatment: last two regimens, if any

Regimen 1: \_\_\_\_\_ Dates/duration of use: \_\_\_\_\_ Response: \_\_\_\_\_

Regimen 2: \_\_\_\_\_ Dates/duration of use: \_\_\_\_\_ Response: \_\_\_\_\_

- Prior partial responder  Prior null responder

Stopped prior therapy for other reason: \_\_\_\_\_

Regimen: \_\_\_\_\_ Dates/duration of use: \_\_\_\_\_ Response: \_\_\_\_\_

### Social History (check all that apply)

- Patient is  $\geq 18$  years of age **OR** meets current AASLD guidelines for treatment

#### Documentation (available if requested) of:

- Counseling regarding abstinence from alcohol, IV drug use and education on how to prevent HCV transmission.
- Abstinence from drugs for at least 6 months; negative urine drug screen required if there is IV drug use history.

#### For women of childbearing potential and male patients with female partners of childbearing potential (for RBV regimens only):

- Patient is not pregnant (or a male with a pregnant female partner) and not planning to become pregnant during treatment or within 6 months of stopping treatment.
- Agreement that partners will use 2 forms of effective contraception during treatment and for at least 6 months after stopping tx.
- Verification that monthly pregnancy tests will be performed throughout treatment.

### Other Medications (OTC, Herbal and Prescription) Information

| Drug name / strength | Frequency / instructions | Quantity | Refills |
|----------------------|--------------------------|----------|---------|
|                      |                          |          |         |
|                      |                          |          |         |

SUBMISSION AND/OR APPROVAL OF A DRUG PRIOR AUTHORIZATION REQUEST DOES NOT GUARANTEE MEDICAID PAYMENT FOR PHARMACY PRODUCTS OR THE AMOUNT OF PAYMENT. ELIGIBILITY FOR AND PAYMENT OF MEDICAID SERVICES ARE SUBJECT TO ALL TERMS AND CONDITIONS AND LIMITATIONS OF THE MEDICAID PROGRAM.

**Confidentiality Notice:** This communication, including any attachments, is for the sole use of the intended recipient(s) and may contain confidential and privileged information. Any unauthorized review, use, disclosure or distribution is prohibited. If you are not the intended recipient, please contact the sender by reply telephone (1-877-537-0722) or fax (1-877-537-0720) and destroy all copies of the original message 1/23/2020

# PRIOR AUTHORIZATION DESCRIPTION

- Note annotations and respective definitions as described on “Hepatitis C Therapy PA Request Guidelines for Regimen Selection and Definitions” page



## Hepatitis C Therapy PA Request Guidelines for Regimen Selection and Definitions

For beneficiaries meeting the criteria, the Mississippi Division of Medicaid (DOM) will approve Hepatitis C treatment PA requests as follows:

- The regimens listed for each clinical scenario below are the preferred regimens for MS Medicaid beneficiaries, based on clinical and cost considerations and are consistent with AASLD/IDSA guidelines.  
<http://www.hcvguidelines.org/full-report-view>

OR

- Pediatric formulations of direct acting antivirals (DAA) with FDA approval will be approved for those patients under the age of eighteen when used according to the table below for treatment naïve children as well as in accordance with current AASLD guidelines including for indication and age - Prior authorization is still required prior to the first dose.**

| GT  | Age (years) | Weight (kg) | Drug                                  | Dose                            | Weeks |
|-----|-------------|-------------|---------------------------------------|---------------------------------|-------|
| Any | 3 to 11     | < 20        | Mavyret Oral Pellets                  | Three 50 mg/20 mg packets daily | 8     |
|     |             | 20 - <30    | Mavyret Oral Pellets                  | Four 50 mg/20 mg packets daily  | 8     |
|     |             | 30 - < 45   | Mavyret Oral Pellets                  | Five 50 mg/20 mg packets daily  | 8     |
|     |             | 45+         | Mavyret Tablets                       | Three 100/40 tablets daily      | 8     |
|     | 12+         |             | Mavyret Tablets                       | Three 100/40 tablets daily      | 8     |
|     | ≥ 6         | ≥ 30        | sofosbuvir/velpatasvir 400/100 tablet | One tablet daily                | 12    |

OR

- Clinical rationale is provided for using a regimen that is beyond those within the current guidelines, or for selecting regimens using non-preferred drugs on the Mississippi Division of Medicaid Universal Preferred Drug List*

On the PA Request Form, which must be approved prior to the 1<sup>st</sup> dose, document the clinical condition supporting the requested regimen. Unless otherwise specified, you do not need to attach documentation to the form. You are attesting to the fact that documentation is available in the patient chart for all information you provide.

### DEFINITIONS/ANNOTATIONS USED ON PA FORM:

▽ **Low Dose Ribavirin = 600 mg/day and increase as tolerated**

◇ **Ribavirin-Ineligible** (documentation exists in the patient’s chart for at least one of the following):

- Hypersensitivity to RBV
- History of severe or unstable cardiac disease
- Pregnant women and men with pregnant partners
- Diagnosis of hemoglobinopathy (e.g., thalassemia major, sickle cell anemia)
- Baseline platelet count < 70,000 cells/mm<sup>3</sup>
- ANC < 1500 cells/mm<sup>3</sup>
- Hb < 12 gm/ml in women or <13 g/dl in men

**RENAL DYSFUNCTION** Patients with CrCl <50 ml/mm should not be treated with ribavirin.

SUBMISSION AND/OR APPROVAL OF A DRUG PRIOR AUTHORIZATION REQUEST DOES NOT GUARANTEE MEDICAID PAYMENT FOR PHARMACY PRODUCTS OR THE AMOUNT OF PAYMENT. ELIGIBILITY FOR AND PAYMENT OF MEDICAID SERVICES ARE SUBJECT TO ALL TERMS AND CONDITIONS AND LIMITATIONS OF THE MEDICAID PROGRAM.

**Confidentiality Notice:** This communication, including any attachments, is for the sole use of the intended recipient(s) and may contain confidential and privileged information. Any unauthorized review, use, disclosure or distribution is prohibited. If you are not the intended recipient, please contact the sender by reply telephone (1-877-537-0722) or fax (1-877-537-0720) and destroy all copies of the original message.  
06/1/2021

# PRIOR AUTHORIZATION DESCRIPTION

- *Note annotations and respective definitions as described on “Hepatitis C Therapy PA Request Guidelines for Regimen Selection and Definitions” page*



## Preferred Direct Acting Antivirals

Mavyret (glecaprevir/pibrentasvir) 300/120 mg  
sofosbuvir/velpatasvir 400/100 mg

## Pediatric Indicated Direct Acting Antiretrovirals (FDA approved age ranges and indications ONLY)

Mavyret (glecaprevir/pibrentasvir) 400/100 mg  
Mavyret Oral Pellets (glecaprevir/pibrentasvir) 50/20mg  
Epclusa 200/50 mg tablet  
Harvoni (ledipasvir/sofosbuvir) 45/200 mg tablet  
Harvoni (ledipasvir/sofosbuvir) 90/400 mg tablet  
Harvoni 33.75/150 mg pellet pak  
Harvoni 45/200 mg pellet pak  
Sovaldi (sofosbuvir) 200 mg  
sofosbuvir/velpatasvir 400/100 mg

## Non-Preferred Direct Acting Antivirals

Harvoni (ledipasvir/sofosbuvir) 90/400 mg  
Sovaldi (sofosbuvir) 400 mg  
ledipasvir/sofosbuvir 90/400 mg  
Vosevi (sofosbuvir, velpatasvir, and voxilaprevir)  
Epclusa (sofosbuvir/velpatasvir) 400/100 mg

**Preferred (most cost-effective) Regimens Listed Below. Not all available regimens are listed.**

**NOTE: Adult Guidelines have changed substantially; most recommendations are largely genotype non-specific; exceptions are noted in red**

## **PLEASE CHECK REQUESTED REGIMEN**

|   |
|---|
| <b>ADULT: Treatment naïve</b>   |
| <b>No cirrhosis</b> <ul style="list-style-type: none"> <li><input type="checkbox"/> Mavyret 100/40 mg, three (3) tablets daily for 8 weeks (for GT5/6 and/or HIV/HCV co-infection, 12 weeks is recommended)</li> <li><input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg, one tablet daily for 12 weeks</li> </ul>         |
| <b>Compensated cirrhosis, HIV negative</b> <ul style="list-style-type: none"> <li><input type="checkbox"/> Mavyret 100/40 mg, three (3) tablets daily for 8 weeks</li> <li><input type="checkbox"/> sofosbuvir/velpatasvir 400/100, one tablet daily for 12 weeks (for GT3, add weight based RBV if Y93H positive)</li> </ul>     |
| <b>Compensated cirrhosis, HIV positive</b> <ul style="list-style-type: none"> <li><input type="checkbox"/> Mavyret 100/40 mg, three (3) tablets daily for 12 weeks</li> <li><input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg, one tablet daily for 12 weeks (for GT3, add weight based RBV if Y93H positive)</li> </ul> |
| <b>ADULT: Treatment experienced (with or without compensated cirrhosis)</b>   |
| <b>Sofosbuvir-based regimen</b> <ul style="list-style-type: none"> <li><input type="checkbox"/> Mavyret 100/40 mg, three (3) tablets daily for 16 weeks</li> </ul>  |
| <b>NS3/4 protease inhibitor inclusive regimen (e.g. Zepatier)</b> <ul style="list-style-type: none"> <li><input type="checkbox"/> Vosevi 400/100/100 mg, one tablet daily for 12 weeks</li> </ul>   |
| <b>Mavyret</b> <ul style="list-style-type: none"> <li><input type="checkbox"/> Vosevi 400/100/100 mg, one tablet daily for 12 weeks (if compensated cirrhosis, add weight based RBV)</li> </ul>   |
| <b>Vosevi or sofosbuvir + Mavyret</b>   |

SUBMISSION AND/OR APPROVAL OF A DRUG PRIOR AUTHORIZATION REQUEST DOES NOT GUARANTEE MEDICAID PAYMENT FOR PHARMACY PRODUCTS OR THE AMOUNT OF PAYMENT. ELIGIBILITY FOR AND PAYMENT OF MEDICAID SERVICES ARE SUBJECT TO ALL TERMS AND CONDITIONS AND LIMITATIONS OF THE MEDICAID PROGRAM.

**Confidentiality Notice:** This communication, including any attachments, is for the sole use of the intended recipient(s) and may contain confidential and privileged information. Any unauthorized review, use, disclosure or distribution is prohibited. If you are not the intended recipient, please contact the sender by reply telephone (1-877-537-0722) or fax (1-877-537-0720) and destroy all copies of the original message.  
06/1/2021

# PRIOR AUTHORIZATION DESCRIPTION

- *Note annotations and respective definitions as described on "Hepatitis C Therapy PA Request Guidelines for Regimen Selection and Definitions" page*



|  |
|--|
| <input type="checkbox"/> Vosevi 400/100/100 mg, one tablet daily + weight based RBV for 24 weeks   |
| <b>GT 3 only: sofosbuvir/NS5A (e.g. Harvoni)</b>   |
| <input type="checkbox"/> Vosevi 400/100/100 mg, one tablet daily + weight based RBV for 12 weeks   |
| <b>ADULT: Re-infection of Allograft Liver after Transplant</b>   |
| <b>DAA-treatment naïve, no decompensated cirrhosis</b>   |
| <input type="checkbox"/> Mavyret 100/40 mg, three (3) tablets daily for 12 weeks   |
| <input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg, one tablet daily for 12 weeks  |
| <b>DAA-treatment experienced, no decompensated cirrhosis</b>   |
| <input type="checkbox"/> Vosevi 400/100/100 mg, one tablet daily for 12 weeks  |
| <b>IF multiple negative baseline characteristics, consider</b>   |
| <input type="checkbox"/> Vosevi 400/100/100 mg, one tablet daily + low dose RBV for 12 weeks   |
| <b>Treatment naïve, decompensated cirrhosis</b>  |
| <input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg, one tablet daily + low dose RBV for 12 weeks   |
| <b>Treatment experienced, decompensated cirrhosis (Child-Pugh B or C ONLY)</b>   |
| <input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg, one tablet daily + low dose RBV for 24 weeks   |
| <b>ADULT: Decompensated Cirrhosis</b>  |
| <b>No prior sofosbuvir or NS5A failure</b>   |
| <input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg + weight-based RBV daily for 12 weeks (low dose RBV recommended for Child-Pugh class C cirrhosis) |
| <input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg daily for 24 weeks (will be approved only for patients with documented ineligibility for RBV)     |
| <b>Prior sofosbuvir or NS5A failure</b>  |
| <input type="checkbox"/> sofosbuvir/velpatasvir 400/100 mg + weight-based RBV daily for 24 weeks (low dose RBV if Child-Pugh C)                              |
| <b>Other Treatment Regimen requested</b>   |
| <b>Genotype, treatment history, and extent of liver disease:</b><br><hr/> <hr/>  |
| <b>Drug names, doses and durations:</b> _____<br><hr/>   |
| <b>Clinical rationale for selecting regimens other than those outlined above:</b><br><hr/> <hr/>   |

## PRIOR AUTHORIZATION DESCRIPTION

- *Note annotations and respective definitions as described on “Hepatitis C Therapy PA Request Guidelines for Regimen Selection and Definitions” page*



**For unique patient populations with renal impairment or HIV:** please refer to the current AASLD Guidelines for recommended treatments. <http://www.hcvguidelines.org/full-report-view>

**If HIV positive:** Please refer to the current AASLD Guidelines for potential drug interactions and recommended dosing adjustments.

### DRUG INTERACTIONS

Reference: <http://hep-druginteractions.org/> provides clinically useful, reliable, and current evidence-based information on relevant drug interactions with hepatitis medications.